arXiv:1611.05309v1 [math.AG] 16 Nov 2016

A LOWER BOUND FOR THE GONALITY CONJECTURE

WOUTER CASTRYCK

Abstract. For every integer $k \geq 3$ we construct a k-gonal curve C along with a very ample divisor of degree $2g+k-1$ (where g is the genus of C) to which the vanishing statement from the Green-Lazarsfeld gonality conjecture does not apply.

The gonality conjecture due to Green and Lazarsfeld [GL86] states that for C a smooth complex projective curve of genus $g \geq 1$ and gonality $k \geq 2$, and L a globally generated divisor on C of sufficiently large degree, one has the following vanishing criterion in Koszul cohomology:

$$K_{i,1}(C,L) \neq 0 \quad \Leftrightarrow \quad 1 \leq i \leq h^0(C,L) - k - 1. \tag{1}$$

This conjecture was proved two years ago by Ein and Lazarsfeld [EL15], who moreover provided the sufficient lower bound $\deg L \geq g^3$. In the meantime this was improved to $\deg L \geq 4g-3$ by Rathmann [Ra16].

It is likely that this bound can be improved further, although Green and Lazarsfeld already noted [GL86, p. 86] that one at least needs

$$\deg L \geq 2g+k-1 \tag{2}$$

because of the non-vanishing of $K_{g-1,1}(K_C + D)$, for any divisor D of rank 1 and degree k on C. Very recently Farkas and Kemeny [FK16] showed that if C is sufficiently generic inside the moduli space of k-gonal curves of genus g then the bound (2) is sufficient for the vanishing criterion (1) to hold. Moreover they conjectured that this should be true for *all* curves whose genus is large enough when compared to the gonality, a statement referred to as the effective gonality conjecture. Results due to Green [Gr84, Thm. 3.c.1] resp. Teixidor i Bigas [Te07, Prop. 3.8] imply that this is indeed the case for trigonal resp. tetragonal curves of genus $g > 3$ resp. $g > 6$.

In this note we aim for an improved delimitation of the foregoing considerations, through the following result.

Theorem. *For each $k \geq 3$ there exists a curve C of genus $g = k(k-1)/2$ along with a very ample divisor L of degree $2g+k-1$ such that*

$$K_{h^0(C,L)-k,1}(C,L) \neq 0. \tag{3}$$

In particular, the bound (2) *is not sufficient for the gonality conjecture to apply.*

The main conclusions to be drawn are that on the one hand, at most, one can hope to improve Rathmann's bound to $\deg L \geq 2g+k$, and that on the other hand $g > k(k-1)/2$ is a necessary lower bound in the statement of Farkas and

Kemeny's effective gonality conjecture. In particular one observes that the special cases implied by the works of Green and Teixidor i Bigas are sharp.

The construction is short and explicit: we let C be a smooth plane curve of degree $k+1$ carrying two total inflection points. By positioning these points at the coordinate points $(1:0:0)$ and $(0:1:0)$ at infinity, and moving the intersection point of the corresponding tangent lines to the finite coordinate point $(0:0:1)$, we can assume that inside $\mathbb{A}^2$ our curve is defined by a polynomial $f(x,y)$ whose Newton polygon Γ is as follows:



Conversely, every curve C defined by a sufficiently generic polynomial that is supported on the above polygon will do. We let $L = (k+1)D$ where

$$D + (1:0:0) + (0:1:0)$$

is the effective divisor cut out by the line at infinity. Being a smooth plane curve of degree $k+1$, the genus of C equals $g = k(k-1)/2$, while its gonality is k by [Se87, Prop. 3.13]. Finally the degree of L equals $(k+1)(k-1) = 2g+k-1$, as announced.

Identifying the function field of C with the field of fractions of $\mathbf{C}[x,y]/(f(x,y))$, we have the following lemma:

Lemma. *L is a very ample divisor, and the vector space $H^0(C,L)$ admits the monomial basis*

$$\{\, x^i y^j \mid i \geq 1, j \geq 1, i+j \leq k+1 \,\}.$$

In particular $h^0(C,L) = \binom{k+1}{2} = k(k+1)/2$.

Proof. The very-ampleness follows from degree considerations. As for the other statement, it follows from basic principles in toric geometry that $H^0(C,L)$ is generated by all monomials that are supported on Γ; see for example [CDV06, Thm. 2] for a ready-to-use statement[1], where we note that our divisor L is exactly the divisor D_C given there. These monomials are given by

$$\{\, 1 \,\} \;\cup\; \{\, x^i y^j \mid i \geq 1, j \geq 1, i+j \leq k+1 \,\},$$

[1]Strictly spoken that theorem only applies if $f(x,y)$ is non-degenerate with respect to its Newton polygon, in our case meaning that C is non-tangent to the line at infinity, but it holds more generally. (On the other hand, for the purpose of proving our main theorem, adding the non-degeneracy assumption is fine.)

which are linearly dependent because $f(x,y) = 0$. Since $f(x,y)$ has a non-zero constant term we can eliminate 1 from the above set, and the remaining monomials are clearly linearly independent functions on C. □

If one uses the above basis of $H^0(C,L)$ to embed C in

$$\mathbb{P}^{\binom{k+1}{2}-1}$$

then it naturally lands inside the $(k-1)$-fold Veronese surface $\nu_{k-1}(\mathbb{P}^2)$. This implies that we have an injection

$$K_{k(k+1)/2-k,1}\left(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2}(k-1)\right) \hookrightarrow K_{k(k+1)/2-k,1}(C,L) = K_{h^0(C,L)-k,1}(C,L).$$

But it is known that the former space is non-trivial; see e.g. [CCDL16, Thm. 1.8]. From this our theorem follows.

Remark. The curve C, when embedded using L, can be viewed as a hyperplane section of the projectively embedded toric surface X_Γ associated to the lattice polygon Γ. Thanks to the Cohen-Macaulay property of toric surfaces, the graded Betti tables of C and X_Γ are the same. In previous joint work with Cools, Demeyer and Lemmens [CCDL16], we had already observed 'exceptional' vanishing behavior among the linear syzygies in the case of X_Γ (there Γ was denoted as Υ_{k-1}). As it turns out, this percolates to the curve level, thereby saying something meaningful about the gonality conjecture.

Remark. We work over $\mathbf{C}$ because for instance [EL15] does so as well, but the theorem presented above is valid over any algebraically closed field, with the same proof.

Acknowledgements

The author was supported by the European Research Council under the European Community's Seventh Framework Programme (FP7/2007-2013) with ERC Grant Agreement 615722 MOTMELSUM, by research project G093913N of the Research Foundation Flanders (FWO), and by the Labex CEMPI (ANR-11-LABX-0007-01). He wishes to thank Gavril Farkas and Milena Hering for some helpful suggestions and comments.

References

[CCDL16] W. Castryck, F. Cools, J. Demeyer, A. Lemmens, *On graded Betti tables of toric surfaces*, preprint

[CDV06] W. Castryck, J. Denef, F. Vercauteren, *Computing zeta functions of nondegenerate curves*, International Mathematics Research Papers **2006**, pp. 1-57 (2006)

[EL15] L. Ein, R. Lazarsfeld, *The gonality conjecture on syzygies of algebraic curves of large degree*, Publications Mathématiques de l'IHÉS **122**(1), pp. 301-313 (2015)

[FK16] G. Farkas, M. Kemeny, *Linear syzygies of curves with prescribed gonality*, preprint

[Gr84] M. Green, *Koszul cohomology and the geometry of projective varieties*, Journal of Differential Geometry **19**(1), pp. 125-171 (1984)

[GL86] M. Green, R. Lazarsfeld, *On the projective normality of complete linear series on an algebraic curve*, Inventiones Mathematicae **83**(1), pp. 73-90 (1986)
[Ra16] J. Rathmann, *An effective bound for the gonality conjecture*, preprint
[Se87] F. Serrano, *Extension of morphisms defined on a divisor*, Mathematische Annalen **277**(3), pp. 395-413 (1987)
[Te07] M. Teixidor i Bigas, *Syzygies using vector bundles*, Transactions of the AMS **359**(2), pp. 897-908 (2007)

Laboratoire Paul Painlevé, Université de Lille-1
Cité Scientifique, 59655 Villeneuve d'Ascq Cedex, France
Departement Elektrotechniek, KU Leuven and imec
Kasteelpark Arenberg 10/2452, 3001 Leuven, Belgium
wouter.castryck@gmail.com